

FLOROMOX (Eye Drops)

MOXIFLOXACIN HYDROCHLORIDE BP

Compositions:

FLOROMOX Sterile Eye Drops: Each ml contains Moxifloxacin Hydrochloride BP equivalent to Moxifloxacin 5 mg. Floromox Sterile Eye Ointment: Each gram contains Moxifloxacin Hydrochloride BP equivalent to Moxifloxacin 5 mg.

Pharmacology:

Moxifloxacin shows antibacterial activity by inhibiting the bacterial enzyme topoisomerase - II (DNA gyrase) and topoisomerase - IV.

Dosage And Administration:

For Eye Drops: One drop in the affected eye 3 times per day for 7 days. For Eye Ointment: It should be applied thinly and evenly to the affected eye three times a day for the first two days and for the next five days apply two times a day or as advised by the registered physician.

Contraindications:

Moxifloxacin is contraindicated in patients with history of hypersensitivity to Moxifloxacin, to other quinolones or to any of components in the medication.

Warning And Precaution:

Prolonged use may result in overgrowth of non-susceptible organisms, including fungi like with other anti-infectives.

Side Effects:

The most frequently reported ocular side effects were conjunctivitis, decreased visual acuity, dry eye, keratitis, ocular discomfort, ocular hyperemia, ocular pain, ocular pruritus, sub-conjunctival hemorrhage and tearing. These events occurred in approximately 1-6% of patients.

Use in Pregnancy and Lactation:

Moxifloxacin should be used during pregnancy only if the potential benefit justifies the potential risk to the fetus as there are no adequate and well-documented studies in pregnant women. Use in lactation: Moxifloxacin has not been measured in human milk, although it can be presumed to be excreted in human milk. Caution should be exercised when Moxifloxacin eye drops & eye ointment is administered to a lactating mother.

Drug Interaction:

Drug-drug interaction studies have not been conducted with Moxifloxacin Hydrochloride. In vitro studies indicate that Moxifloxacin does not inhibit CYP3A4, CYP2D6, CYP2C9, CYP2C19, or CYP1A2 indicating that Moxifloxacin is unlikely to alter the pharmacokinetics of drugs metabolized by these cytochrome P450 isoenzymes.

Overdosage:

No information available.

Storage:

Store at 2°&-25°& C and dry place, protect from light. Keep out of reach of children.

Packing:

FLOROMOX Eye Drops: Each plastic dropper bottle contains 5 ml sterile eye drops. FLOROMOX Eye Ointment: Each tube contains 5 gm sterile eye ointment.

Manufactured By:
The IBN SINA Pharmaceutical Industry PLC.
Shafipur, Gazipur, Bangladesh.