

# **ANODYNE (Eye Drops)**

DICLOFENAC SODIUM BP

## **Compositions:**

Each ml contains Diclofenac Sodium BP 1 mg.

## **Pharmacology:**

Diclofenac Sodium having analgesic & anti inflammatory action. Diclofenac inhibits the cyclooxygenase enzyme essential for the biosynthesis of prostaglandin which is responsible for inflammation.

## **Dosage And Administration:**

Pre-operative: Use upto 5 drops over 3 hours preceding surgery. Post operative: Use 3 drops immediately after surgery, then 3 to 5 drops daily for as long as required. Others: Use 4 to 5 drops daily, depending on the severity of the condition. Children: No-specific investigation has been carried out.

## **Contraindications:**

Diclofenac eye drops should not be used in patient with known hypersensitivity to Diclofenac or any of its excipients. Like other NSAID, Diclofenac is contraindicated in patients suffering from asthma, urticaria or acute rhinitis.

## **Warning And Precaution:**

Diclofenac eye drops should be used only after removal of contact lens in patients who use contact lens. If infection is present, appropriate treatment with antibiotic should be given concomitantly. The drug should not be prescribed during 3rd trimester of pregnancy.

## **Side Effects:**

Occasional mild to moderate transient stinging and blurred vision may occur after application of the Diclofenac eye drops. Hypersensitivity reaction in the form of itching, reddening and photosensitivity occur rarely.

## **Use in Pregnancy and Lactation:**

During pregnancy, Diclofenac should be employed only for compelling reasons. The lowest effective dose should be used. These types of drugs are not recommended during the first trimester of pregnancy. In view of insufficient clinical data, Diclofenac Sodium Gel is not recommended during pregnancy. A very insignificant quantity of Diclofenac may be detected in breast milk but no undesirable effects on the infant to be expected.

## **Drug Interaction:**

No information available

## **Overdosage:**

No information available

## **Storage:**

Store in a cool, dry place and protected from light. Keep out of reach of children. Discard the container 4 weeks after opening.

## **Packing:**

Anodyne & Eye Drops: Each plastic dropper bottle contains 5ml sterile eye drops.

Manufactured By:  
The IBN SINA Pharmaceutical Industry PLC.  
Shafipur, Gazipur, Bangladesh.